

## Astaxanthin inhibits cytokines production and inflammatory gene expression by suppressing I $\kappa$ B kinase-dependent nuclear factor $\kappa$ B activation in pre and postpartum Murrah buffaloes during different seasons

Lakshmi Priyadarshini and Anjali Aggarwal

Animal Physiology Division, ICAR-National Dairy Research Institute, Karnal - 132 001, Haryana, India.

**Corresponding author:** Lakshmi Priyadarshini, e-mail: drlpaa@gmail.com

**Co-author:** AA: anjaliaggarwal23aa@gmail.com

**Received:** 23-12-2017, **Accepted:** 04-05-2018, **Published online:** 10-06-2018

**doi:** 10.14202/vetworld.2018.782-788 **How to cite this article:** Priyadarshini L, Aggarwal A (2018) Astaxanthin inhibits cytokines production and inflammatory gene expression by suppressing I $\kappa$ B kinase-dependent nuclear factor  $\kappa$ B activation in pre and postpartum Murrah buffaloes during different seasons, *Veterinary World*, 11(6): 782-788.

### Abstract

**Aim:** We examined regulatory function of astaxanthin on mRNA expression of nuclear factor  $\kappa$ B (NF- $\kappa$ B) p65, interleukin-6 (IL-6), tumor necrosis factor alpha (TNF- $\alpha$ ), and interferon gamma (IFN- $\gamma$ ) in peripheral blood mononuclear cells in pre and postpartum Murrah buffaloes during summer (temperature-humidity index [THI]=86; relative humidity [RH]=24) and winter (THI=58.74; RH=73) seasons.

**Materials and Methods:** A total of 32 Murrah buffaloes apparently healthy and in their one to four parity were selected from National Dairy Research Institute herd and equally distributed randomly into four groups (control and supplemented groups of buffaloes during summer and winter season, respectively). All groups were fed according to the nutrient requirement of buffaloes (ICAR, 2013). The treatment group was supplemented with astaxanthin at 0.25 mg/kg body weight/animal/day during the period 30 days before expected date of calving and up to 30 days postpartum.

**Results:** There was downregulation of NF- $\kappa$ B p65 gene in all the groups. NF- $\kappa$ B p65 mRNA expression was lower ( $p < 0.05$ ) in treatment than control group from prepartum to postpartum during summer, while mRNA expression was low only on day 21 after calving during winter season. The mRNA expression of IL-6, TNF- $\alpha$ , and IFN- $\gamma$  was lower ( $p < 0.05$ ) in treatment than a control group of buffaloes during summer and winter seasons. The mRNA expression of NF- $\kappa$ B p65, IL-6, TNF- $\alpha$ , and IFN- $\gamma$  was higher ( $p < 0.05$ ) in summer than in winter seasons.

**Conclusion:** The xanthophyll carotenoid astaxanthin a reddish-colored C-40 compound is a powerful broad-ranging antioxidant that naturally occurs in a wide variety of living organisms, such as microalgae, fungi, crustaceans, and complex plants. Astaxanthin blocked nuclear translocation of NF- $\kappa$ B p65 subunit and I $\kappa$ B $\alpha$  degradation, which correlated with its inhibitory effect on I $\kappa$ B kinase (IKK) activity. These results suggest that astaxanthin, probably due to its antioxidant activity, inhibits the production of inflammatory mediators by blocking NF- $\kappa$ B activation and as a consequent suppression of IKK activity and I $\kappa$ B-degradation.

**Keywords:** astaxanthin, Murrah buffalo, nuclear factor  $\kappa$ B p65, summer, winter.

### Introduction

Astaxanthin is a lipophilic, pinkish-orange carotenoid (3,3'-dihydroxy- $\beta$ , $\beta$ -carotene-4,4'-di-one) found in algae, seafood (crustacean shells, crab, shrimps, and fish) by Qin *et al.* [1] and various plants, giving them their exclusive colored aspect [2]. The main source of AST is the microalga *Haematococcus pluvialis*, which contains maximum concentrations [3]. Astaxanthin is also used as a dietary additive in the USA, Japan, South Korea, and Sweden [4]. Like another carotenoid, astaxanthin manifests high protective antioxidant [5,6] and anticancer [7,8] properties which decrease oxidative stress and inflammation [9,10],

reduces rethrombosis after thrombolysis [11] and is efficient in ischemia-reperfusion [12], arterial hypertension [13], and dyslipidemia [14].

Nuclear factor  $\kappa$ B (NF- $\kappa$ B) is a well-characterized transcription factor that is known to regulate the expression of a wide range of genes, including those of cytokines, chemokines, and cytokine receptors [15] that control various aspects of the immune [16] and inflammatory response [17]. The liberated NF- $\kappa$ B translocates to the nucleus and binds as a transcription factor to  $\kappa$ B motifs in the promoters of target genes, leading to their transcription. Aberrant NF- $\kappa$ B activity is associated with various inflammatory diseases, and most anti-inflammatory drugs suppress inflammatory cytokine expression by inhibiting the NF- $\kappa$ B pathway [18,19]. Thus, an NF- $\kappa$ B inhibitor has clinical potential in inflammatory diseases. NF- $\kappa$ B responds to oxidative stress promptly and regulates the transcription of genes involved in a wide range of antioxidant, immune and inflammatory cell functions [20]. NF- $\kappa$ B activation is highly dependent on oxidative stress,

Copyright: Priyadarshini and Aggarwal. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

which is associated with the immune response [21,22]. An imbalance in oxidants/antioxidants, an excess of oxidants and/or a depletion of antioxidants, can lead to oxidative stress [23]. The inhibition or clearance of intracellular reactive oxygen species generation and accumulation by nicotinamide adenine dinucleotide phosphate oxidase decreased NF- $\kappa$ B activation [24].

We hypothesized that astaxanthin regulated the production of pro-inflammatory cytokines by inhibiting NF- $\kappa$ B activation during different seasons. We studied effect of astaxanthin (at 0.25 mg/kg body weight/animal) as an antioxidant on mRNA expression of NF- $\kappa$ B p65, IL-6, TNF- $\alpha$ , and interferon gamma (IFN- $\gamma$ ) in peripheral blood mononuclear cells (PBMCs) in pre and postpartum Murrah buffaloes during summer and winter seasons.

## Materials and Methods

### Ethical approval

The experiment was approved by the Institutional Animal Ethics Committee (approval no. 41-IAEC-18-10) constituted as per the article 13 of the CPCSEA rules, laid down by Government of India. All the ethical guidelines were followed during the course of the experiment.

### Chemicals

Histopaque 1077 from Sigma Chemical Co. (St. Louis, MO, USA) and Dulbecco's phosphate buffer saline (DPBS) procured from HiMedia Laboratories Pvt. Ltd., India was used for separation of PBMCs. RNeasy (Cat. No-74104) and RNase-Free DNase Set (Cat No. - 79251) were purchased from QIAGEN Pvt. Ltd., New Delhi, India. Nuclease-free water was purchased from Genetix, Biotech Asia Pvt. Ltd (Fermentas life sciences). Revert Aid First strand cDNA synthesis kit (K1622, Thermo Scientific), Dream Taq Green PCR Master Mix (2X) (K 1082), DyNAmo Color Flash SYBER Green Qpcr kit (F-416XL), and GeneRuler™ 100 bp DNA Ladder (50  $\mu$ g) were purchased from Thermo Fisher Scientific (LSG) Mumbai, India.

### Experimental animals

A total of 32 animals, eight each of pregnant buffalo (*Bubalus bubalis*) having an approximate body weight of 600-650 kg, milk yield of (average) 7.6 kg in their one to four parity 30 days before parturition from Livestock Research Center (LRC), National Dairy Research Institute (NDRI), Karnal, were selected for the experiment. The experiments were carried out during two distinct phases coinciding with two seasons of the year, namely winter (December-February) and summer (April-June). From these 32 animals were further randomly divided equally (8 each) into four groups (control and supplemented groups of buffaloes during summer and winter seasons, respectively). The experimental animals were given adaptation period of 1 week to get adapted to new surroundings. Control groups of buffaloes during summer and winter seasons remained without provision of astaxanthin

supplementation, while treatment groups of buffaloes were managed with astaxanthin supplementation during summer and winter seasons, respectively. The astaxanthin as powder form was fed at 0.25 mg/kg body weight/day [25] mixing with concentrate mixture, from 30 days before parturition till 30 days after parturition. At the time of experiment, all the animals were clinically healthy and free from any abnormalities. The experimental animals were maintained and fed as per standard practices followed at LRC, NDRI, and Karnal.

### Blood collection and RNA extraction

About 6-7 ml of fresh blood samples was drawn aseptically from each animal in potassium-EDTA coated Vacutainer tubes (BD-PlymouthPL6 7BP, UK) and was immediately transported to laboratory under refrigeration. Blood samples were collected at 30, 21, 15, and 7 days prepartum, day of calving and 7, 15, and 21 days postpartum from each animal in both seasons.

### Isolation of PBMC

Bovine PBMCs were isolated after blood centrifugation at 3000 rpm at 4°C for 30 min in a refrigerated centrifuge (Sigma, Germany) for separation of lymphocytes. The buffy coat was harvested and re-suspended in 1:1 v/v DPBS. PBMCs were isolated by density gradient centrifugation method using lymphocyte separation medium, Histopaque 1077 (Sigma). The whole content was layered carefully onto the Histopaque to produce a clean interface between the two layers. Further, it was centrifuged at 2000 rpm for 30 min at room temperature. The white opaque mononuclear fraction from the interface was collected between the DPBS and Histopaque. Further, centrifugation was done at 1500 rpm for 7 min for washing the cells with PBS (pH=7.4). Finally, the cell pellet was obtained.

### RNA extraction

Total RNA from PBMC was isolated using RNeasy Mini Kit (Qiagen India Pvt. Ltd.) according to the manufacturer's protocol. The quality and integrity of isolated RNA was checked by carrying out agarose gel electrophoresis in 1.5% agarose in 1 $\times$  TAE buffer at 100 Volts for 30 min. The RNA purity was verified by optical density (OD) absorption ratio at  $\lambda$ 260/ $\lambda$ 280 using Biospec-nano Spectrophotometer (Shimadzu Corp., Japan). A ratio of ~2.0 was generally accepted as "pure" for RNA. The RNA samples with good purity and integrity were used for cDNA synthesis.

### cDNA synthesis

For each sample, about 200 ng of total RNA was used for cDNA synthesis using Revert Aid First strand cDNA synthesis kit (Fermentas, USA) by reverse transcription polymerase chain reaction (PCR) according to the manufacturer's protocol. The RT reaction was carried out at 25°C for 10 min, 42°C for 60 min, and 75°C for 5 min in a thermal cycler (Bio-Rad, USA).

### Primers

Primers for NF- $\kappa$ B, IL-6, TNF- $\alpha$ , and INF- $\gamma$  were designed using primer 3.0 software or suggested from published literature. The sequences and expected PCR product length are shown in Table-1 and Figure-1.

### Quantitative real-time polymerase chain reaction (RT-PCR)

The RT-PCR reaction was carried out in Applied Biosystems 7500 RT-PCR systems using 0.5  $\mu$ l of cDNA, 5  $\mu$ l of Maxima SYBR green qPCR master mix, and 0.5  $\mu$ l of concerned gene sequence-specific forward and reverse primers (10 pmol), and the final volume of 10  $\mu$ l was made with nuclease-free water. The RT-PCR program consisted of initial heating at 50°C for 2 min followed by 95°C for 10 min, and samples were amplified for 40 cycles (95°C for 30 s; 58°C for NF- $\kappa$ B, IL-6, TNF- $\alpha$  and INF- $\gamma$ , 59°C for GAPDH for 30 s, and 72°C for 30 s). The final extension at 72°C incubation was continued for a further 10 min GAPDH was used as housekeeping gene. The relative

quantification of target gene was done by the  $2^{-\Delta\Delta CT}$  method [26].

### Statistical analysis

The data were analyzed using SAS Software, version 9.3 of the SAS System, Copyright<sup>©</sup> (2011) SAS Institute Inc., Cary, NC, USA. Data from different experiments are presented as mean $\pm$ standard error. The pairwise comparison was drawn using Tukey's multiple comparison tests. The difference at  $p < 0.05$  was considered to be statistically significant. The model performed for each test is as follows:


$$Y_{ij} = \mu + \alpha_i + \beta_j + H_k + e_{ij}$$

Where,  $Y_{ij}$  is a dependent variable,  $\mu$  is the overall mean,  $\alpha_i$  effect of buffaloes ( $i = 1, 2, 3, \dots, 16$ );  $\beta_j$  = effect of treatment  $j$  ( $j$ =astaxanthin supplemented or non-supplemented);  $H_k$  = effect of time period and  $e_{ij}$  is the random error which is assumed to be independent and normally distributed.

### Results

During summer season, the NF- $\kappa$ B p65 gene expression of control and treatment groups was down-regulated significantly ( $p < 0.05$ ) on day of calving and on day 21 after calving as compared to day 30 before calving, respectively. The NF- $\kappa$ Bp65 gene expression decreased significantly ( $p < 0.05$ ) in treatment group as compared to control group on day 7 before calving, on day of calving and during postpartum period. During winter season, the NF- $\kappa$ Bp65 gene expression of control and treatment groups was downregulated significantly ( $p < 0.05$ ) on day of calving and on day 21 after calving as compared to day 30 before calving, respectively. The NF- $\kappa$ B p65 gene expression decreased significantly ( $p < 0.05$ ) in treatment groups as compared to control group on day 60 after calving (Table-2). Gene expression of NF- $\kappa$ B p65 was positively correlated to genes IL-6, TNF $\alpha$ , and INF- $\gamma$  (Table-3). The NF- $\kappa$ Bp65 gene expression was higher ( $p < 0.05$ ) in summer than in winter season.

During the summer season, the IL-6 gene expression in control and treatment groups increased significantly ( $p < 0.05$ ) on the day of calving and day 21 after calving as compared to day 30 before calving, respectively. The IL-6 gene expression decreased significantly ( $p < 0.05$ ) in treatment groups as compared to control group of respective days. During the



**Figure-1:** Agarose gel electrophoresis of Real time PCR amplified products of (a) NF $\kappa$ B p65 (192 bp); (b) IL-6 (105 bp); (c) TNF- $\alpha$  (130 bp) and (d) IFN- $\gamma$  (120 bp) gene products on 1.5% agarose.

**Table-1:** Sequence, annealing temperature, and fragment size of the pairs of primers used for RT-PCR.

Gene	Primer sequence	Annealing temp. (°C)	Fragment size (bp)	Accession no.
NF- $\kappa$ B p65 (RELA)	F-TCCAAGTTCCCATAGAAGAGCA R-TCCAGAGTTCCGATTCACC	57	192	DQ355511
IL-6	F-ATGACGAGTGTGAAAGCAGC R-TCGCCTGATTGAACCCAGAT	59	105	AY347710.1
TNF- $\alpha$	F-GGTCAACATCCTGTCTGCCA R-ACTGAGGCGATCTCCCTTCT	59	130	XM_006041930.1
Interferon $\gamma$	F-CGCAAAGCCATAAATGAACCTC R-GGACCATTACGTTGATGCTC	60	120	AF484688.1

PCR=Polymerase chain reaction

**Table-2:** Mean ( $\pm$ SE) mRNA expression of NF- $\kappa$ B p65 gene in control and treatment groups of buffaloes during pre and postpartum period in different seasons.

Days relative to calving	Summer		Winter	
	Control	Treatment	Control	Treatment
-30	1.00 <sup>dA</sup> $\pm$ 0.00	1.00 <sup>fA</sup> $\pm$ 0.00	1.00 <sup>gA</sup> $\pm$ 0.00	1.00 <sup>fA</sup> $\pm$ 0.00
-21	0.23 <sup>aA</sup> $\pm$ 0.03	0.21 <sup>aA</sup> $\pm$ 0.03	0.19 <sup>aA</sup> $\pm$ 0.01	0.13 <sup>aA</sup> $\pm$ 0.01
-15	0.61 <sup>bA</sup> $\pm$ 0.02	0.67 <sup>cdA</sup> $\pm$ 0.03	0.52 <sup>bA</sup> $\pm$ 0.02	0.45 <sup>bA</sup> $\pm$ 0.03
-7	0.67 <sup>bA</sup> $\pm$ 0.03	0.52 <sup>bb</sup> $\pm$ 0.02	0.67 <sup>cdA</sup> $\pm$ 0.03	0.62 <sup>cA</sup> $\pm$ 0.02
0	0.91 <sup>cdA</sup> $\pm$ 0.04	0.74 <sup>deB</sup> $\pm$ 0.02	0.71 <sup>deA</sup> $\pm$ 0.02	0.69 <sup>cdA</sup> $\pm$ 0.04
7	0.98 <sup>dA</sup> $\pm$ 0.03	0.78 <sup>eB</sup> $\pm$ 0.02	0.78 <sup>eA</sup> $\pm$ 0.02	0.72 <sup>dA</sup> $\pm$ 0.02
15	1.22 <sup>eA</sup> $\pm$ 0.03	0.91 <sup>fB</sup> $\pm$ 0.04	0.89 <sup>fA</sup> $\pm$ 0.05	0.83 <sup>eA</sup> $\pm$ 0.02
21	0.82 <sup>cA</sup> $\pm$ 0.02	0.63 <sup>cB</sup> $\pm$ 0.02	0.61 <sup>bcA</sup> $\pm$ 0.02	0.51 <sup>bb</sup> $\pm$ 0.03

<sup>a,b,c,d,e,f,g</sup> and <sup>A,B</sup> means with different superscripts differ significantly ( $p < 0.05$ ) in a column and in a row of each season, respectively. NF- $\kappa$ B=Nuclear factor  $\kappa$ B, SE=Standard error

**Table-3:** Correlation among various genes.

	NF- $\kappa$ B p65	IL-6	TNF- $\alpha$	Interferon $\gamma$
NF- $\kappa$ B p65	1			
IL-6	0.304**	1		
TNF- $\alpha$	0.346**	0.687**	1	
Interferon $\gamma$	0.695**	0.536**	0.630**	1

\*\* $p < 0.01$ . NF- $\kappa$ B=Nuclear factor  $\kappa$ B, IL-6=Interleukin-6, TNF- $\alpha$ =Tumor necrosis factor alpha

winter season, the IL-6 gene expression in control and treatment groups increased significantly ( $p < 0.05$ ) on the day of calving and day 21 after calving as compared to day 30 before calving, respectively. The IL-6 gene expression decreased significantly ( $p < 0.05$ ) in treatment groups as compared to control group on day 15, 7 before calving and on day 15, 21 after calving (Table-4). The IL-6 gene expression was higher ( $p < 0.05$ ) in summer than in winter season. IL-6 gene expression was positively correlated to genes NF- $\kappa$ B p65, TNF $\alpha$ , and IFN $\gamma$  (Table-3).

During the summer season, the TNF- $\alpha$  gene expression in control and treatment groups increased significantly ( $p < 0.05$ ) on the day of calving and day 21 after calving as compared to day 30 before calving, respectively. The TNF- $\alpha$  gene expression was decreased significantly ( $p < 0.05$ ) in the treatment group as compared to control group. During the winter season, the TNF- $\alpha$  gene expression in control and treatment groups increased significantly ( $p < 0.05$ ) on the day of calving and day 21 after calving as compared to day 30 before calving, respectively. The TNF- $\alpha$  gene expression decreased significantly ( $p < 0.05$ ) in the treatment group as compared to control group on day 15<sup>th</sup> before calving to day 15<sup>th</sup> after calving, respectively (Table-5). The TNF- $\alpha$  expression was higher ( $P < 0.05$ ) in summer than in winter season. TNF- $\alpha$  gene expression was positively correlated to genes NF- $\kappa$ B p65, IL-6, IFN- $\gamma$ .

During the summer season, the IFN- $\gamma$  gene expression in control and treatment groups was upregulated significantly ( $p < 0.05$ ) from day of calving to day 15 then downregulated significantly ( $p < 0.05$ ) on day 21 after calving as compared to day 30 before calving, respectively. The IFN- $\gamma$  gene expression decreased

significantly ( $p < 0.05$ ) in treatment groups as compared to control group from day 7 before calving to day 21 after calving. During the winter season, the IFN- $\gamma$  gene expression of control and treatment groups was significantly ( $p < 0.05$ ) upregulated on day of calving to day 15 then downregulated significantly ( $p < 0.05$ ) on day 21 after calving as compared to day 30 before calving, respectively (Table-6). The IFN- $\gamma$  gene expression decreased significantly ( $p < 0.05$ ) in treatment groups as compared to control group from the day of calving to day 15 after calving. The IFN- $\gamma$  gene expression was higher ( $p < 0.05$ ) in summer than winter season. IFN- $\gamma$  gene expression was positively correlated to genes NF- $\kappa$ B p65, TNF $\alpha$ , and IL-6 (Table-3).

## Discussion

NF- $\kappa$ B p65 is one of the most important pathways in inflammation and tumors [27]. In our experiment, the mRNA expression of NF- $\kappa$ B p65, IL-6, TNF $\alpha$ , and IFN- $\gamma$  significantly ( $p < 0.05$ ) decreased in astaxanthin supplemented groups than that of control groups in summer and winter seasons. Astaxanthin decreased the expression of pro-inflammatory mediators such as TNF- $\alpha$  by suppressing I $\kappa$ B-dependent NF- $\kappa$ B activation in both primary macrophages and lipopolysaccharide (LPS)-stimulated RAW264.7 cells [28]. Astaxanthin remarkably suppressed the expression of the mRNA expression of inflammatory mediators, i.e., tumor necrosis factor alpha (TNF- $\alpha$ ), IL-6, and in THP-1 macrophages [29]. Astaxanthin significantly decreased the production of IL-6 and TNF- $\alpha$  in LPS-stimulated neutrophils [30]. Astaxanthin treatment reduced the secretion of IL-6 and TNF- $\alpha$  in H<sub>2</sub>O<sub>2</sub>-stimulated U937 mononuclear cells, and this property was elicited by a restoration of the basal SHP-1 protein expression level and reduced NF- $\kappa$ B (p65) nuclear expression [31]. Another study reported that on macrophage NF- $\kappa$ B activation inhibition might be due to antioxidant property of astaxanthin [32]. IL-6 and TNF- $\alpha$  gene expression decreased ( $p < 0.05$ ) after farrowing and in lactation [33]. IFN- $\gamma$  expression upregulated significantly ( $p < 0.05$ ) from day of calving to day 15 then after was downregulated significantly ( $p < 0.05$ ) on day 21 after calving as compared to day

**Table-4:** Mean ( $\pm$ SE) mRNA expression of IL-6 gene in control and treatment groups of buffaloes during pre and postpartum period in different seasons.

Days relative to calving	Summer		Winter	
	Control	Treatment	Control	Treatment
-30	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00
-21	1.08 <sup>aA</sup> $\pm$ 0.03	1.03 <sup>aA</sup> $\pm$ 0.02	1.04 <sup>aA</sup> $\pm$ 0.02	1.01 <sup>aA</sup> $\pm$ 0.01
-15	1.43 <sup>bA</sup> $\pm$ 0.03	1.32 <sup>bA</sup> $\pm$ 0.03	1.28 <sup>bA</sup> $\pm$ 0.02	1.15 <sup>bB</sup> $\pm$ 0.02
-7	2.71 <sup>eA</sup> $\pm$ 0.02	2.14 <sup>eB</sup> $\pm$ 0.02	2.06 <sup>dA</sup> $\pm$ 0.03	1.93 <sup>dB</sup> $\pm$ 0.03
0	4.13 <sup>fA</sup> $\pm$ 0.02	3.64 <sup>gB</sup> $\pm$ 0.03	3.32 <sup>fA</sup> $\pm$ 0.02	3.21 <sup>fA</sup> $\pm$ 0.03
7	2.68 <sup>eA</sup> $\pm$ 0.02	2.48 <sup>fB</sup> $\pm$ 0.03	2.21 <sup>eA</sup> $\pm$ 0.02	2.10 <sup>eA</sup> $\pm$ 0.02
15	2.16 <sup>dA</sup> $\pm$ 0.03	2.02 <sup>dB</sup> $\pm$ 0.03	2.03 <sup>dA</sup> $\pm$ 0.03	1.89 <sup>dB</sup> $\pm$ 0.03
21	1.82 <sup>cA</sup> $\pm$ 0.03	1.69 <sup>cA</sup> $\pm$ 0.03	1.79 <sup>cA</sup> $\pm$ 0.03	1.65 <sup>cB</sup> $\pm$ 0.03

<sup>a,b,c,d,e,f,g</sup> and <sup>A,B</sup> means with different superscripts differ significantly ( $p < 0.05$ ) in a column and in a row of each season, respectively. IL-6=Interleukin-6, SE=Standard error

**Table-5:** Mean ( $\pm$ SE) mRNA expression of TNF- $\alpha$  gene in control and treatment groups of buffaloes during pre and postpartum period in different seasons.

Days relative to calving	Summer		Winter	
	Control	Treatment	Control	Treatment
-30	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00	1.00 <sup>aA</sup> $\pm$ 0.00
-21	1.13 <sup>bA</sup> $\pm$ 0.02	1.10 <sup>aA</sup> $\pm$ 0.03	1.08 <sup>aA</sup> $\pm$ 0.03	1.03 <sup>aA</sup> $\pm$ 0.02
-15	2.43 <sup>dA</sup> $\pm$ 0.03	1.83 <sup>bB</sup> $\pm$ 0.03	1.70 <sup>bA</sup> $\pm$ 0.03	1.51 <sup>bB</sup> $\pm$ 0.03
-7	2.78 <sup>eA</sup> $\pm$ 0.02	1.97 <sup>cB</sup> $\pm$ 0.03	1.95 <sup>dA</sup> $\pm$ 0.03	1.67 <sup>cB</sup> $\pm$ 0.03
0	3.12 <sup>fA</sup> $\pm$ 0.02	2.85 <sup>dB</sup> $\pm$ 0.02	2.10 <sup>eA</sup> $\pm$ 0.03	1.93 <sup>dB</sup> $\pm$ 0.03
7	4.91 <sup>gA</sup> $\pm$ 0.02	3.71 <sup>eB</sup> $\pm$ 0.03	3.14 <sup>fA</sup> $\pm$ 0.03	2.92 <sup>eB</sup> $\pm$ 0.02
15	2.37 <sup>dA</sup> $\pm$ 0.03	2.03 <sup>cB</sup> $\pm$ 0.03	2.15 <sup>eA</sup> $\pm$ 0.03	1.91 <sup>dB</sup> $\pm$ 0.03
21	2.10 <sup>cA</sup> $\pm$ 0.03	1.74 <sup>bB</sup> $\pm$ 0.03	1.54 <sup>bA</sup> $\pm$ 0.03	1.41 <sup>bA</sup> $\pm$ 0.03

<sup>a,b,c,d,e</sup> and <sup>A,B</sup> means with different superscripts differ significantly ( $p < 0.05$ ) in a column and in a row of each season, respectively. TNF- $\alpha$ =Tumor necrosis factor alpha, SE=Standard error

**Table-6:** Mean ( $\pm$ SE) mRNA expression of interferon  $\gamma$  gene in control and treatment groups of buffaloes during pre and postpartum period in different seasons.

Days relative to calving	Summer		Winter	
	Control	Treatment	Control	Treatment
-30	1.00 <sup>dA</sup> $\pm$ 0.00	1.00 <sup>eA</sup> $\pm$ 0.00	1.00 <sup>eA</sup> $\pm$ 0.00	1.00 <sup>eA</sup> $\pm$ 0.00
-21	0.07 <sup>aA</sup> $\pm$ 0.02	0.04 <sup>aA</sup> $\pm$ 0.01	0.09 <sup>aA</sup> $\pm$ 0.02	0.02 <sup>aA</sup> $\pm$ 0.00
-15	0.31 <sup>bA</sup> $\pm$ 0.05	0.28 <sup>bA</sup> $\pm$ 0.04	0.26 <sup>bA</sup> $\pm$ 0.04	0.14 <sup>bA</sup> $\pm$ 0.04
-7	0.78 <sup>cA</sup> $\pm$ 0.07	0.52 <sup>cB</sup> $\pm$ 0.02	0.51 <sup>cA</sup> $\pm$ 0.03	0.43 <sup>cA</sup> $\pm$ 0.03
0	1.91 <sup>eA</sup> $\pm$ 0.03	1.64 <sup>fB</sup> $\pm$ 0.03	1.71 <sup>fA</sup> $\pm$ 0.03	1.33 <sup>fB</sup> $\pm$ 0.03
7	2.71 <sup>fA</sup> $\pm$ 0.02	2.03 <sup>gB</sup> $\pm$ 0.03	2.62 <sup>gA</sup> $\pm$ 0.06	2.06 <sup>gB</sup> $\pm$ 0.04
15	2.92 <sup>gA</sup> $\pm$ 0.02	2.15 <sup>hB</sup> $\pm$ 0.02	2.83 <sup>hA</sup> $\pm$ 0.02	2.62 <sup>hB</sup> $\pm$ 0.06
21	0.91 <sup>cdA</sup> $\pm$ 0.04	0.71 <sup>dB</sup> $\pm$ 0.02	0.73 <sup>dA</sup> $\pm$ 0.07	0.65 <sup>dA</sup> $\pm$ 0.02

<sup>a,b,c,d,e,f,g,h</sup> and <sup>A,B</sup> means with different superscripts differ significantly ( $p < 0.05$ ) in a column and in a row of each season, respectively. SE=Standard error

30 before calving. A study on rat found that astaxanthin inhibits inducible nitric oxide synthase enzyme activity, which decreases the production of nitric oxide as well as prostaglandin E2 and TNF- $\alpha$  [34].

IFN- $\gamma$  expression increased gradually from prepartum (0.56-fold) up to 2<sup>nd</sup> week (2.89-fold) postpartum and subsequently decreased in the 3<sup>rd</sup> (0.73-fold) and 4<sup>th</sup> week (0.70-fold) in healthy buffaloes [35]. IL-6 and IFN- $\gamma$  mRNA expressions in liver were higher in astaxanthin fed male broiler chickens (*Gallus gallus*) compared to that of control birds [36].

### Conclusion

We showed that NF- $\kappa$ B inhibited *in vivo* production of pro-inflammatory cytokines, such as TNF- $\alpha$ ,

IL-6, and IFN- $\gamma$ . Furthermore, it suppressed NF- $\kappa$ B activation may be by inhibiting IKK activity. These results support the idea that astaxanthin prevents inflammatory processes by blocking the expression of pro-inflammatory genes as a consequence of suppressing NF $\kappa$ B activation.

### Authors' Contributions

LP carried out the research under the supervision of AA. LP collected blood samples, did molecular work and analyzed the data. The manuscript was drafted and revised by LP under the guidance of AA. Both authors read and approved the final manuscript.

## Acknowledgments

This study was financially supported by National Initiative on Climate Resilient Agriculture, Indian Council of Agricultural Research (NICRA-ICAR), New Delhi (Grant no. 2049/3033). The authors gratefully acknowledge Director, ICAR-National Dairy Research Institute, Karnal, for providing necessary facilities to conduct this study.

## Competing Interests

The authors declare that they have no competing interests.

## References

- Qin, S., Liu, G.X. and Hu, Z.Y. (2008) The accumulation and metabolism of astaxanthin in *Scenedesmus obliquus*. *Process Biochem.*, 43: 795-802.
- Zhao, Z.W., Cai, W., Lin, Y.L., Lin, Q.F., Jiang, Q., Lin, Z. and Chen, L.L. (2011) Ameliorative effect of astaxanthin on endothelial dysfunction in streptozotocin-induced diabetes in male rats. *Arzneimittelforschung*, 61: 239-246.
- Boussiba, S., Cohen, Z., Richmond, A. and Vonshak, A. (2000) Procedure for Large-Scale Production of Astaxanthin from *Haematococcus*. PCT Patents 9, 728, 274.
- Choi, H.D., Kang, H.E., Yang, S.H., Lee, M.G. and Shin, W.G. (2011) Pharmacokinetics and first-pass metabolism of astaxanthin in rats. *Br. J. Nutr.*, 105: 220-227.
- Aoi, W., Naito, Y., Takamami, Y., Ishii, T., Kawai, Y., Akagiri, S., Kato, Y., Osawa, T. and Yoshikawa, T. (2008) Astaxanthin improves muscle lipid metabolism in exercise via inhibitory effect of oxidative CPTI modification. *Biochem. Biophys. Res. Commun.*, 366: 892-897.
- Derosa, G., Limas, C.P., Macias, P.C., Estrella, A. and Maffioli, P. (2014) Dietary and nutraceutical approach to Type 2 diabetes. *Arch. Med. Sci.*, 10: 336-344.
- Kavitha, K., Kowshik, J., Kishore, T.K., Baba, A.B. and Nagini, S. (2013) Astaxanthin inhibits NF-kappaB and Wnt/beta-catenin signaling pathways via inactivation of Erk/MAPK and PI3K/Akt to induce intrinsic apoptosis in a hamster model of oral cancer. *Biochem. Biophys. Acta*, 1830: 4433-4444.
- Palazzo, L., Thomas, B., Jemth, A.S., Coldy, T., Leidecker, O., Feijs, K.L.H., Zaja, R., Loseva, O., Puigvert, J.C., Matic, I., Helleday, T. and Ahel, I. (2015) Processing of protein ADP-ribosylation by nudix hydrolases. *Biochem. J.*, 468(2): 293-301.
- Wolf, A.M., Asoh, S., Hiranuma, H., Ohsawa, I., Iio, K., Satou, A., Ishikura, M. and Ohta, S. (2010) Astaxanthin protects mitochondrial redox state and functional integrity against oxidative stress. *J. Nutr. Biochem.*, 21(5): 381-389.
- Sahebkar, A., Serban, C., Ursosiu, S. and Banach, M. (2015) Effect of curcuminoids on oxidative stress: A systematic review and meta-analysis of randomized controlled trials. *J. Funct. Foods*, 18: 898-909.
- Lauver, D.A., Driscoll, E.M. and Lucchesi, B.R. (2008) Disodium disuccinate astaxanthin prevents carotid artery rethrombosis and *ex vivo* platelet activation. *Pharmacology*, 82: 67-73.
- Gross, G.J. and Lockwood, S.F. (2004) Cardioprotection and myocardial salvage by a disodium disuccinate astaxanthin derivative. *Life Sci.*, 75: 215-224.
- Hussein, G., Goto, H., Oda, S., Sankawa, U., Matsumoto, K. and Watanabe, H. (2006) Antihypertensive potential and mechanism of action of astaxanthin: III. Antioxidant and histopathological effects in spontaneously hypertensive rats. *Biol. Pharm. Bull.*, 29: 684-688.
- Banach, M., Serban, C., Aronow, W.S., Rvsz, J., Dragan, S., Lerma, E.V., Apetrii, M. and Covic, A. (2014) Lipid, blood pressure and kidney update. *Int. Urol. Nephrol.*, 46: 947-961.
- Baeuerle, P.A. and Baichwal, V.R. (1997) NF-kappa B as a frequent target for immunosuppressive and anti-inflammatory molecules. *Adv. Immunol.*, 65: 111-137.
- Dale, E., Davis, M. and Faustman, D.L. (2006) A role for transcription factor NF-kappaB in autoimmunity: Possible interactions of genes, sex, and the immune response. *Adv. Physiol. Educ.*, 30: 152-158.
- Li, N. and Karin, M. (1999) Is NF-kB the sensor of oxidative stress? *FASEB J.*, 13: 1137-1143.
- Castrillo, A., de Las, H.B., Hortelano, S., Rodriguez, B., Villar, A. and Bosca, L. (2001) Inhibition of the nuclear factor kappa B (NF-kB) pathway by tetracyclic kaurene diterpenes in macrophages. Specific effects on NF-kB-inducing kinase activity and on the coordinate activation of ERK and p38 MAPK. *J. Biol. Chem.*, 276: 15854-15860.
- Keifer, J.A., Guttridge, D.C., Ashburner, B.P. and Baldwin, A.S. Jr. (2001) Inhibition of NF-kB activity by thalidomide through suppression of Ikb kinase activity. *J. Biol. Chem.*, 276: 22382-22387.
- Ghosh, S. and Hayden, M.S. (2008) New regulators of NF-kB in inflammation. *Nat. Rev. Immunol.*, 8: 837-848.
- Romagnoli, M., Gomez-Cabrera, M.C., Perrelli, M.G., Biasi, F., Pallardó, F.V., Sastre, J., Poli, G. and Vina, J. (2010) Xanthine oxidase-induced oxidative stress causes activation of NF-kappaB and inflammation in the liver of type I diabetic rats. *Free Radic. Biol. Med.*, 49(2): 171-177.
- Hybertson, B.M., Gao, B., Bose, S.K. and McCord, J.M. (2011) Oxidative stress in health and disease: The therapeutic potential of Nrf2 activation. *Mol. Aspects. Med.*, 32: 234-246.
- Pedernera, M., Celi, P., Garcia, S.C., Salvin, H.E., Barchia, I. and Fulkerson, W.J. (2010) Effect of diet, energy balance and milk production on oxidative stress in early-lactating dairy cows grazing pasture. *Vet. J.*, 186: 352-357.
- Kim, H., Na, H.J., Kim, C.K., Kim, Y., Ha, K.S., Lee, H., Chung, H.T., Kwon, H.J., Kwon, Y.G. and Kim, Y.M. (2008) The non-provitamin a carotenoid, lutein, inhibits NF-kB dependent gene expression through redox-based regulation of the phosphatidylinositol 3 kinase/PTEN/Akt and NF-kB-inducing kinase pathways: Role of H<sub>2</sub>O<sub>2</sub> in NF-kB activation. *Free Radic. Biol. Med.*, 45: 885-896.
- Lignell, A. and Inborr, J. Method of the Prophylactic Treatment of Mastitis. United State Patent US 6335015 B1, January 1, 2002.
- Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup>. *Method.*, 25(4): 402-408.
- Li, D., Song, J.Z., Li, H., Shan, M.H., Liang, Y., Zhu, J. and Xie, Z. (2015) Storage lipid synthesis is necessary for autophagy induced by nitrogen starvation. *FEBS Lett.*, 589(2): 269-276.
- Lee, S.J., Bai, S.K., Lee, K.S., Namkoong, S., Na, H.J., Ha, K.S., Han, J.A., Yim, S.V., Chang, K., Kwon, Y.G., Lee, S.K. and Kim, Y.M. (2003) Astaxanthin inhibits nitric oxide production and inflammatory gene expression by suppressing I-kappaB kinase-dependent NF-kappaB activation. *Mol. Cells*, 16: 97-105.
- Kishimoto, Y., Tani, M., Uto-Kondo, H., Iizuka, M., Saita, E., Sone, H., Kurata, H. and Kondo, K. (2010) Astaxanthin suppresses scavenger receptor expression and matrix metalloproteinase activity in macrophages. *Eur. J. Nutr.*, 49: 119-126.
- Macedo, R.C., Bolin, A.P., Marin, D.P. and Otton, R. (2010) Astaxanthin addition improves human neutrophils function: *In vitro* study. *Eur. J. Nutr.*, 49: 447-457.
- Speranza, L., Pesce, M., Patrino, A., Franceschelli, S., de Lutiis, M.A., Grilli, A. and Felaco, M. (2012) Astaxanthin treatment reduced oxidative induced pro-inflammatory cytokines secretion in U937: SHP-1 as a novel biological target. *Mar. Drugs*, 10(4): 890-899.
- Kishimoto, Y., Yoshida, H. and Kondo, K. (2016) Potential

- anti-atherosclerotic properties of astaxanthin. *Mar. Drugs*, 14(35): 1-13.
33. Contreras, G.A., Kirkwood, R.N. and Sordillo, L.M. (2013) Mononuclear leukocyte fatty acid composition and inflammatory phenotype in periparturient and lactating sows. *J. Anim. Sci.*, 91: 174-187.
34. Ohgami, K., Shiratori, K., Kotake, S., Nishida, T., Mizuki, N., Yazawa, K. and Ohno, S. (2003) Effects of astaxanthin on lipopolysaccharide-induced inflammation *in vitro* and *in vivo*. *Invest. Ophthalmol. Vis. Sci.*, 44(6): 2694-2701.
35. Patra, M.K., Kumar, H. and Nandi, S. (2013) Neutrophil functions and cytokines expression profile in buffaloes with impending postpartum reproductive disorders. *Asian Aust. J. Anim. Sci.*, 26(10): 1406-1415.
36. Takahashi, K., Takimoto, T., Sato, K. and Akiba, Y. (2011) Effect of dietary supplementation of astaxanthin from *Phaffiarhodozyma* on lipopolysaccharide-induced early inflammatory responses in male broiler chickens (*Gallus gallus*) fed a corn-enriched diet. *Anim. Sci. J.*, 82(6): 753-758.

\*\*\*\*\*