

Windrow composting as an option for disposal and utilization of dead birds

G. Vinodkumar¹, V. Ramesh Saravanakumar¹, S. Ramakrishnan², A. Elango³ and S. C. Edwin⁴

1. Department of Livestock Production and Management, Veterinary College and Research Institute, Namakkal- 637 002, Tamil Nadu, India; 2. Veterinary University Training and Research Centre, Tiruvannmalai - 606601, Tamil Nadu, India; 3. Department of Dairy Science, Veterinary College and Research Institute, Namakkal - 637 002, Tamil Nadu, India; 4. Department of Livestock Production and Management, Veterinary College and Research Institute, Tirunelveli- 627001, Tamil Nadu, India.

Corresponding author: G. Vinodkumar, GV: drvinodlpm@gmail.com, VRS: lpmramesh1@yahoo.com, SR: ramkilpnamakkal@rediffmail.com, AE: elango@tanuvas.org.in, SCE: drscedwin@tanuvas.org.in

Received: 04-03-2014, **Revised:** 19-04-2014, **Accepted:** 23-04-2014, **Published online:** 06-06-2014

doi: 10.14202/vetworld.2014.377-379

How to cite this article: Vinodkumar G, Saravanakumar VR, Ramakrishnan S, Elango A and Edwin SC (2014) Windrow composting as an option for disposal and utilization of dead birds, *Veterinary World* 7(6): 377-379.

Abstract

Aim: The present study was undertaken to ascertain the feasibility of windrow composting as an environmentally safe and bio-secure disposal method of poultry manure and mortalities.

Materials and Methods: Poultry dead birds and cage layer manure were collected from the commercial poultry farms and coir pith was obtained from coir fiber extraction unit. Physical properties and chemical composition of ingredients were analyzed and a suitable compost recipe was formulated. Two treatment windrow groups (T₁- Dead birds + Cage layer manure + Coir pith, T₂- Cage layer manure + Coir pith) in replication were fabricated. Physical chemical and biological parameters of compost were analyzed.

Results: Temperature profile ensured maximum pathogen and parasite reduction. Reduction in moisture content, weight, volume, total organic carbon, and progressive increase in total ash, calcium, phosphorus and potassium content as the composting proceeded, were indicative of organic matter degradation and mineralization. Favourable C:N ratio and germination index indicated compost maturity and absence of any phytotoxins in finished compost. The finished compost had undetectable level of *Salmonella*. There was no odour and fly menace throughout the composting experiment.

Conclusion: Windrow composting of poultry waste can be considered as a biologically and environmentally safe disposal option with recycling of nutrients in the form of compost.

Keywords: dead birds, poultry, windrow composting.

Introduction

The poultry industry in India has undergone a paradigm shift, from backyard venture before the 1960s to vibrant intensive agribusiness, growing at around 8 to 10% annually over the last decade. This fast growing industry is currently facing issues regarding waste management. Accumulation of large amount of wastes is one of the major problems, posing serious challenge to the further development of industry [1]. Poultry industry waste includes excreta, mortalities, feathers, hatchery waste, waste feed and broken eggs. These are largely organic materials and are convertible to useful resources. It has become challenging task for poultry farmers to handle these enormous quantities of waste, especially mortalities. Current methods for routine disposal of poultry mortalities include burial, incineration and anaerobic digestion in closed pits [2]. Each one of these have their own inherent disadvantages like cost involved, labour intensiveness, production of environmental pollutants and obnoxious odour etc.

Composting is one of the most versatile and

remunerative techniques for handling such biodegradable solid wastes by biological stabilization into a safer and more stabilized material called as compost [3], which can be used as a source of nutrients and soil conditioner in agricultural applications [4].

Hence the present study was undertaken to know the feasibility of windrow composting as environmentally safe and biosecure method of disposal option for poultry manure and mortalities.

Materials and Methods

A composting experiment was carried out at Veterinary College and Research Institute, Namakkal, Tamil Nadu, India, during 2013. The study included poultry wastes, such as dead birds and cage layer manure and coir pith as co-composting material. Poultry dead birds and cage layer manure were collected from the commercial poultry farms and coir pith was obtained from coir fiber extraction unit. Physical properties and chemical composition of these were analyzed and a compost recipe was formulated with suitable C:N ratio (20:1) and moisture content of and 60%. Windrows were fabricated on impervious flooring with the alternative layers of mixture of fresh cage layer manure and coir pith and dead birds (T₁). Quantity of ingredients *viz.*, dead birds, cage layer manure and coir pith

Copyright: The authors. This article is an open access article licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>) which permits unrestricted use, distribution and reproduction in any medium, provided the work is properly cited.

Table-1: Mean \pm SE of physical and chemical properties of poultry waste compost

Parameters	Primary stage			Secondary stage		
	T ₁	T ₂	'F' value	T ₁	T ₂	'F' value
Peak temperature (°C)	65.70 \pm 0.10	57.50 \pm 0.30	-	62.30 \pm 0.00	49.35 \pm 0.85	-
Persistence of thermophilic (55 °C and above) temperature (days)	18.50 \pm 0.50	04.50 \pm 0.00	-	05.50 \pm 1.50	00.00 \pm 0.00	-
Composting period (days)	34.50 \pm 0.50	20.00 \pm 1.00	-	23.50 \pm 2.50	09.50 \pm 0.50	-
Moisture content (%)	30.52 \pm 1.41	27.98 \pm 1.01	NS	33.87 \pm 0.92	30.68 \pm 0.69	*
Weight reduction (%)	-	-	-	21.08 \pm 0.56	26.40 \pm 0.70	-
Volume reduction (%)	-	-	-	14.13 \pm 1.55	15.68 \pm 1.58	-
pH	7.06 \pm 0.06	8.32 \pm 0.10	**	7.58 \pm 0.11	7.84 \pm 0.13	NS
EC in mS/cm	4.54 \pm 0.09	4.74 \pm 0.12	NS	3.68 \pm 0.08	3.86 \pm 0.06	NS
Total ash (%)	58.90 \pm 0.93	58.33 \pm 0.35	NS	60.78 \pm 0.58	62.30 \pm 0.30	*
Total carbon (%)	23.84 \pm 0.54	24.17 \pm 0.21	NS	22.75 \pm 0.34	21.87 \pm 0.17	*
Total nitrogen (g/kg)	13.44 \pm 0.43	11.67 \pm 0.60	*	13.52 \pm 0.39	11.71 \pm 0.48	**
C:N ratio	17.91 \pm 0.60	23.57 \pm 1.34	**	16.98 \pm 0.56	21.32 \pm 0.82	**
Calcium, (g/kg)	32.47 \pm 1.44	38.03 \pm 1.29	**	59.94 \pm 0.96	58.58 \pm 0.60	NS
Phosphorus, (g/kg)	19.69 \pm 1.80	21.21 \pm 0.69	NS	22.16 \pm 0.27	22.23 \pm 0.47	NS
Potassium, (g/kg)	30.69 \pm 0.92	32.27 \pm 1.00	NS	32.06 \pm 1.11	33.94 \pm 0.71	NS

** , Significant at 1% level $P < 0.01$; * , Significant at 5% level $P < 0.05$; NS, Non significant

was in the ratio of 1:3:2.4 based on the compost recipe formulated. Another windrow consisting of mixture of cage layer manure and coir pith in the ratio of 4.4:1 (T₂) was constructed. Moisture level in these was adjusted to 60 per cent by addition of water. Composting process was carried out as per the procedure followed by USDA-NRCS.

Physical analysis: Windrow temperature was monitored daily, using digital thermometer with long probe. Sufficient quantity of finished compost samples from each windrow from different locations were collected in an air tight polythene bag and transferred immediately for moisture estimation. Moisture content of composting samples was determined as weight loss upon drying at 105 °C in the hot air oven for 24 hours. Electrical conductivity (EC) and pH were measured using water proof multi parameter pocket tester (EUTECH Instrument) by preparing 1: 10 w/v compost – water extract.

Chemical analysis: Total organic matter (TOM) was calculated by gravimetric loss on ignition produced by ashing the samples in a muffle furnace for 24 hours at 430 °C. The total organic carbon content was calculated from the ash content using the formula:

$$\text{Total organic carbon} = [1 - \text{ash content} \times (1000)].$$

The total carbon was calculated from total organic matter value using the conventional 'Van Bemelem factor' of 1.724. The weight loss on ignition was divided by 1.724 to give the percentage of total carbon. Compost samples were analysed for total Kjeldahl nitrogen, phosphorus and calcium. The concentration of potassium was analyzed using Flame photometer as described by Jackson [5]. The data thus collected were statistically analyzed as per the methods suggested by Snedecor and Cochran [6].

Biological parameters: Compost maturity was evaluated by germination index of cow pea and sorghum [7]. Presence of *Salmonella* was tested in finished compost by enumerating the microbes on *Salmonella Shigella* agar [8].

Results and Discussion

Mean \pm SE of different physical and chemical parameters of poultry waste compost samples at the primary and secondary stage of composting are given in Table-1. Core temperature of both windrows began to rise swiftly and reached thermophilic temperature within three days, indicating favorable conditions like C:N ratio, moisture, aeration and availability of degradable organic matter for microbial activity [9, 10]. Windrow's core peak temperature and persistence of thermophilic temperature in windrows with dead birds were higher in windrows with dead birds. Persistence of thermophilic temperature *ie.* above 55 °C for minimum period of three consecutive days in poultry mortality composting will kill parasites and fecal and plant pathogens within the pile [10]. In this experiment all windrows had fairly higher than standard minimum period of persistence of thermophilic temperature and ensured the bio-safety. Such prolonged period of persistence of thermophilic temperature in dead bird composting was also recorded previously [11-15].

Composting of dead birds along with cage layer manure was completed within 58 days, which was within the range of earlier reports ranging widely between 14 and 127.5 days [1, 11-14, 16, 17, 18]. Due to microbial degradation of organic matter and subsequent heat generation, there was reduction in moisture, weight and volume of windrows at the end of composting [3, 19]. Finished compost of dead birds had slightly alkaline pH suggesting stabilization and was in line with earlier reports with pH ranging from 6.82 to 8.9 [11-14, 18].

Total ash, calcium, phosphorus and potassium content progressively increased and total organic content decreased as composting proceeded, indicating mineralization and degradation of organic matter [20]. Significant reduction in nitrogen was observed in windrows due to ammonia volatilization [19]. The C:N ratio in windrows with dead birds reduced below 20:1 indicating the maturity [3]. The minimum germination index of finished compost recorded in this experiment

was 85.54% indicating that compost samples were free from phytotoxic substances [7]. The finished compost had undetectable level of *Salmonella*. There was no odour and fly menace throughout the composting experiment.

Conclusion

The results indicated that windrow composting of poultry wastes viz., dead birds and cage layer manure with coir pith as co-composting material can be considered as biologically and environmentally safe disposal option, with recycling nutrients in the form of compost.

Author's contributions

VRS, SR, AE and SCE designed work plan, GV conducted experiment under supervision of VRS, SR, AE, SCE. Laboratory analysis of samples was carried out by GV under the guidelines of VRS and AE. VRS did statistical analysis and critically reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors are thankful to the Dean, Veterinary College and Research Institute, Namakkal, Tamil Nadu, India for providing the necessary facilities to carry out the work. The authors are thankful to authority of Tamil Nadu Veterinary Animal Sciences University, India for providing fund for this study.

Competing interests

The authors declare that they have no competing interests.

References

- Kelleher, B.P., Leahy, J.J., Henihan, M., O'Dwyer, T.F., Sutton, D. and Leahy, M.J. (2002) Advances in poultry litter disposal technology--a review. *Bioresour. Technol.*, 83(1): 27-36.
- Kumar, V.R.S., Sivakumar K., Purushothaman M.R., Natarajan A. and Amanullah, M.M., (2007) Chemical Changes During Composting of Dead Birds with Caged Layer Manure. *J. Appl. Sci. Res.*, 3(10): 1100-1104.
- Gajalakshmi, S., and Abbasi, S.A. (2008) Solid Waste Management by Composting: State of the Art. *Crit. Rev. Env. Sci. Technol.*, 38: 311-400.
- Gabhane, J., William, S.P.M.P., Bidyadhar, R., Bhilawe, P., Anand, D., Vaidya, A.N., and Wate, S.R. (2012) Additives aided composting of green waste: effects on organic matter degradation, compost maturity, and quality of the finished compost. *Bioresour. Technol.*, 114: 382-8.
- Jackson, M.L. (1973) Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi. p224.
- Snedecor, G.W. and Cochran, W.G. (1989) Statistical Methods. 8th Edition, Iowa State Press, Ames, Iowa, USA., p254-268.
- Tiquia, S.M., Tam, N.F.Y. and Hodgkiss, I.J. (1996) Microbial activities during composting of spent pig-manure sawdust litter at different moisture contents. *Bioresour. Technol.*, 55:201-206.
- Sivakumar, K., (2006) Disposal and utilisation of poultry carcasses by aerobic composting. PhD Thesis submitted to Tamil Nadu Veterinary and Animal science University Chennai, India.
- Gu, W., Zhang F., Xu, P., Tang, S., Xie, K., Huang, X. and Huang, Q. (2011) Bioresource Technology Effects of Sulphur and Thiobacillus Thioparus on Cow Manure Aerobic Composting. *Bioresour. Technol.*, 102(11): 6529-6535.
- Nelson, V. (2011) Poultry mortality composting. Agdex 450/29-1, Livestock mortality documents, Alberta Agriculture and Rural Development, Canada. p1-36.
- McCaskey, T.A. 1994. Dead bird composting . Final report for Contract USDA-43-2D81-1-561, Auburn University, p27.
- Costa, M.S.S., de M., Costa, L.A.de M., Pelá, A., da Silva, C.J., Decarli, L.D. and Matter, U.F. (2006) Desempenho de quatro sistemas para compostagem de carcaça de aves. *R. Bras. Eng. Agríc. Ambiental*, 10(3):692-698.
- Sakthivadivu, R. (2008) Composting of poultry waste using low cost carbonaceous materials. MVSc Thesis submitted to Tamil Nadu Veterinary and Animal science University Chennai, India.
- Lawson, M.J. and Keeling A.A. (1999) Production and physical characteristics of composted poultry carcasses. *Br. Poult. Sci.*, 40: 706-708.
- Paiva, E. C. R., Matos, A.T., Azevedo, M.A. Barros, R. and Costa, T. (2012) Evaluation of composting poultry carcasses by the methods of composting windrows and aerated static. *Eng Agric.* 32 (5):1-6.
- Collins, E.R., (1996). Composting Dead Poultry. Fact sheet, Virginia cooperative extension, Virginia State University http://www2.ca.uky.edu/poultryprofitability/Production_manual/Chapter17_Dead_bird_disposal/Chapter17_Composting_dead_poultry.pdf. Accessed on 12-05-2013.
- Abreu, P.G., Paiva, D.P., Abreu, V.M.N., Coldebella, A. and Cestonaro, T. (2011) Casca de arroz e palhada de soja na compostagem de carcaças de frangos de corte. *Acta Scientiarum. Anim. Sci.*, 33(1): 51-57.
- Ahmed, Z.A.M, Hussin, H.A., Rohaim, M.A. and Nsar, S.A.E. (2012) Efficacy of Composting Dead Poultry and Farms Wastes Infected with Avian Influenza Virus H5N1. *Am. Euras. J. Agric. & Environ. Sci.*, 12(5): 588-96.
- Bertoldi, M.D., Vallini, G. And Pera, A. (1983). The biology of composting. *Waste Manage. Res.* 1: 157-176.
- Chefetz, B., Hatcher, P.G., Hadar, Y. and Chen, Y. (1996). Chemical and biological characterization of organic matter during composting of municipal solid waste. *J. Environ. Qual.*, 25: 776-785.
