

Study on inclusion of probiotic, prebiotic and its combination in broiler diet and their effect on carcass characteristics and economics of commercial broilers

M. A. Saiyed¹, R. S. Joshi², F. P. Savaliya³, A. B. Patel³, R. K. Mishra³ and N. J. Bhagora³

1. Veterinary Dispensary Tarapur, District Panchayat Anand, Gujarat, India; 2. Department of Animal Genetics & Breeding, Veterinary College, Anand Agricultural University, Anand, Gujarat, India; 3. Poultry Complex, Veterinary College, Anand Agricultural University, Anand, Gujarat, India.

Corresponding author: M. A. Saiyed, email: dr.shahid92@gmail.com, RSJ: rsjoshi106@gmail.com, FPS: fpsavaliya@yahoo.com, ABP: patelab1411@yahoo.co.in, RKM: rkmishra@gmail.com, NJB: nikeshbhagora@gmail.com
Received: 09-12-2014, **Revised:** 03-01-2015, **Accepted:** 09-01-2015, **Published online:** 25-02-2015

doi: 10.14202/vetworld.2015.225-231. **How to cite this article:** Saiyed MA, Joshi RS, Savaliya FP, Patel AB, Mishra RK, Bhagora NJ (2015) Study on inclusion of probiotic, prebiotic and its combination in broiler diet and their effect on carcass characteristics and economics of commercial broilers, *Veterinary World* 8(2):225-231.

Abstract

Background and Aim: In today era, broiler industry facing a problem of price hiking of feed of broiler, also in competitive era there should be lower feed cost, lower feed conversion ratio, low feed consumption yet good body weight at marketable age.

Materials and Methods: Day-old commercial broiler chicks (n=200) were distributed randomly into 5 dietary treatment groups viz. control (T₁), probiotic in the feed @ 100 g/tonne of feed (T₂), prebiotic in the feed @ 500 g/tonne of feed (T₃), probiotic + prebiotic @ 100 g/tonne and 500 g/tonne of feed, respectively (T₄) and probiotic + prebiotic @ 50 g/tonne and 250 g/tonne of feed (T₅). The growth of broilers and dressing weight along with the weight of giblet (liver without gall bladder, gizzard without serous layer, and heart without pericardium), Kidney, Abdominal fat, Length of Intestine and dressing percentage were measured. Economics in terms of Return Over Feed Cost (ROFC) and European Performance Efficiency Index (EPEI) was calculated.

Results: Among all carcass traits, dressing percentage, abdominal fat weight and abdominal fat percentage (as a percentage of dressed weight) were recorded significant (p<0.05) difference among different treatment groups. The income from selling of the birds was significantly (p<0.05) higher in all treatment groups than the control group but there was a non-significant difference between supplemented groups. Feed cost during whole experimental period was significantly (p<0.05) lower in synbiotic supplemented groups (T₄ and T₅) than other groups. ROFC of all treatment group found significantly (p<0.05) higher than the control group.

Conclusion: It can be concluded that the diet supplemented with synbiotic (100% level) was most efficient in terms of EPEI and synbiotic (50% level) in terms of ROFC. Hence, as feed supplement, synbiotic has a beneficial effect over probiotic and prebiotic when used alone.

Keywords: broiler, carcass traits, European Performance Efficiency Index, prebiotic, probiotic, Return Over Feed Cost, symbiotic.

Introduction

Poultry is one of the fastest growing segments among the component of livestock sector in India. Production of agricultural crops has been growing at a rate of 1.5-2.0% per annum while poultry industry is growing at 8-15% per annum in India. The organized sector of the poultry is contributing nearly 70% of the total output and the rest 30% by the unorganized sector. About 66.7% of the total output from poultry is realized from the poultry meat sector and only 33.3% from egg production. The total poultry population, which was only 73.5 million in 1951 has made tremendous growth during the past 50 years and has reached 648.83 million as per 2007 Census. India occupied the 3rd position in egg production and 5th position in poultry meat production as per 2010 census in the

world. The per capita availability of poultry meat is 2.15 kg/annum, which is very less as against the recommendation of 11 kg meat/annum given by National Institute of Nutrition [1].

Feed as a major input item to broiler rearing for being 75% of the production cost has a vital role in broiler economics. Hence, it is imperative to give due attention to proper utilization of feed without adversely affecting the growth or production performance of broilers [2]. Antibiotics have been also used to promote growth rate, improve feed conversion ratio (FCR) and reduce mortality in broiler flocks. However, repeated use of antibiotics in poultry diets resulted in severe problems like resistance of pathogen to antibiotics, accumulation of antibiotics residue in their products and environment, imbalance of normal microflora and reduction in beneficial intestinal microflora [3,4].

This has led to the development of different products to be used as feed additives such as enzymes, probiotics, prebiotics, organic acids, and plant extracts. Probiotics are "live microorganisms, which when

Copyright: The authors. This article is an open access article licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>) which permits unrestricted use, distribution and reproduction in any medium, provided the work is properly cited.

administered in adequate amounts confer a health benefit on the host" [5]. Whereas prebiotics are defined as non-digestible food ingredients that beneficially affect the host by selectively stimulating the growth and activity of one or a limited number of bacteria. Synbiotics refer to nutritional supplements combining probiotics and prebiotics and in a form of synergism. The main reason, for using a synbiotic, is that a true probiotic, without its prebiotic food, does not survive well in the digestive system. Without the necessary food source for the probiotic, it will have a greater intolerance for oxygen, low pH, and temperature [6]. Poultry meat production has been paid more and more attention because it is particularly high in quantities and qualities of valuable protein, essential amino acids, fat, essential fatty acids, vitamins and minerals. Poultry meat is a high quality concentrated food and, therefore, plays an important role in human nutrition [7]. Poultry meat constitutes around 20% of total meat production in the country [8].

Materials and Methods

Ethical approval

The present study was carried out at the age of 42 days/6 week and carcass parameters were carried out at chicken shop with due care after selling of the birds.

Experimental design

A 200-day-old commercial broiler chicks were randomly distributed into five different treatment groups having four replicates in each treatment with 10 chicks in each replicate and they were reared under battery brooder system of cage up to 42 days of age.

Experimental procedure

Feed and water were offered *ad libitum*, and standard managerial practices followed. Chicks were weighed individually at the start of the experiment and at the end of every week. Mortality was also recorded. Composition of dietary treatments is given in Table-1. Proportion of different feed ingredient is shown in Table-2 and detailed composition of feed additives is shown in Table-2.1 First group of birds were kept as a control, and they were not supplied feed with either probiotic or prebiotic in both, broiler starter and broiler finisher feed. Probiotic in the feed of T₂ group was given at the rate of 100 g/tonne of feed during the starter phase (0-4 weeks) and finisher

phase (5-6 weeks). Prebiotic in the feed of T₃ group was given at the rate of 500 g/tonne of feed during the starter phase and finisher phase. In T₄ group, probiotic was given at the rate of 100 g/tonne of feed and Prebiotic was given at the rate of 500 g/tonne of feed during the starter phase and finisher phase.

Table-2: Proportion of feed ingredients (%) used for preparation of broiler starter and finisher feeds.

Ingredients	Proportion (Kg)	
	Broiler starter	Broiler finisher
Maize	53.900	54.075
Deoiled rice bran	1.150	9.150
Soyabean DOC	39.755	30.500
Trace mineral ¹	0.100	0.000
Shell grit	2.200	1.700
DCP	2.000	1.390
Salt	0.300	0.400
Enzymes ²	0.050	0.050
Furazolidone ³	0.025	0.050
Metabolic activator ⁴	0.100	0.100
Toxin binder ⁵	0.100	0.100
Maduramicin ⁶	0.050	0.050
Lysine	0.100	0.100
DLMethionine	0.135	0.090
Herbal performance enhancer ⁷	0.025	0.025
Vitamin B ₁₂ ⁸	0.010	0.020
Vegetable oil	0.000	2.000
Vitamin and mineral supplement ⁹	0.000	0.200
Total	100.00	100.00
CP (%) [*]	23.10	20.20
ME (kcal/kg feed) [*]	2800.00	2900.00

*As per calculated values, DCP=Digestible crude protein

Table-2.1: Detailed composition of feed additives added in feed.

- Trace Minerals:
Each kg contains:
Copper-15 g, Iodine-1 g, Iron-60 g, Manganese-80 g, Selenium-0.3 g, Zinc-80 g, Inorganic nutritive care-Q.S.
- Enzymes:
Each gram contains:
Xylanase-2000 IU, Amylase-400 IU, Protease-4000 IU, Cellulase-500 IU.
- Furazolidone:
Each kg contains:
Furazolidone-200 g, Inorganic nutritive carrier-Q.S.
- Metabolic activator:
Lecithin extract treated with co enzyme.
- Toxin binder:
Selected silicates, surfactants, organic acids and salts of organic acids.
- Maduramicin:
Ionophore (polyether antibiotic) coccidiostat.
- Herbal performance enhancer @ 250 g per tonne of feed.
- Vitamin B₁₂:
Each kg contains: Vitamin B₁₂-100 mg
- Vitamin and mineral supplement:
Each 2 kg contains:
Vitamin A-50 lakh IU, Vitamin B2-2 g, Vitamin B6-400 mg, Vitamin B12-5600 mcg, Vitamin E-800 IU, Iron-7.5 g, Vitamin D3-6.25 lakh IU, Choline chloride-10 g, Copper-2 g, Iodine-1 g, Zinc-15 g, Manganese-27.5 g, Calcium-27.25%, Phosphorus-7.45%, Calcium pantothenate-4 g.

Table-1: Composition of dietary treatment employed.

Dietary treatment	Composition
T ₁	Basal diet without additive, served as control
T ₂	Basal diet+Probiotic @ 100 g/tonne of feed
T ₃	Basal diet+Prebiotic @ 500 g/tonne of feed
T ₄	Basal diet+Probiotic @ 100 g/tonne of feed+Prebiotic @ 500 g/tonne of feed
T ₅	Basal diet+Probiotic @ 50 g/tonne of feed+Prebiotic @ 250 g/tonne of feed

Whereas, in T₅ group, probiotic was given at the rate of 50 g/tonne of feed and prebiotic was given at the rate of 250 g/tonne of feed during starter phase and finisher phase i.e. half of the dose than T₄ treatment. Each gram of probiotic contains 6 × 10⁹ colony forming units of *Lactobacillus acidophilus*, *Lactobacillus casei*, *Pediococcus acidilactici*, *Bacillus subtilis* and *Saccharomyces boulardii*. Whereas, prebiotic contains Mannan Oligo-Saccharide in which Mannan 12-14% and Glucan 13-16% were included. The broiler starter (0-28 days) and broiler finisher (29-42 days) feeds for different treatments were prepared as per the guidelines of BIS (1992). Return Over Feed Cost (ROFC) is calculated by subtracting the feed cost during the rearing period from the income from the sold bird on live weight basis. Whereas European Performance Efficiency Index (EPEI) (European Performance Efficiency Index), as described by [9], was calculated by following formula.

$$\text{EPEI} = \frac{\text{Ave. body weight (g)} \times \text{Livability (\%)} \div 10}{\text{Age of flock (days)} \times \text{FCR}}$$

Statistical analysis

The data on various traits were analyzed using completely randomized design as per Snedecor and Cochran (1995) [10].

Results and Discussion

The carcass traits like dressed weight, dressing%, liver weight, heart weight, gizzard weight, giblet weight, giblet%, abdominal fat weight, abdominal fat%, intestinal length and kidney weight of the different groups offered feed additives either singly or in combination in relation to control is presented in Table-3.

Dressed weight

The average body weight gain for different treatment group and for weekly interval as well as for starter and finisher phase is given in Table-4. The highest dressed weight was observed in synbiotic half level group (T₅), which was followed by T₂, T₃, T₄ and T₁ groups. Dressed weight was highest in T₅ though it had the lowest pre-slaughter weight. The dressed weight differences were non-significant amongst all treatment groups. Present findings were in accordance with [11-14].

Dressing yield (%)

The highest dressing per cent was observed in synbiotic half level group (T₅) which was followed by T₃, T₂, T₄ and T₁ group and all treatment groups dressing percentages were significantly (p<0.05) higher than control group but with non-significant difference between supplemental groups. Present findings differed from [11-13,15,16] as they all observed non-significant differences between supplemental groups and

Table-3: Carcass characteristics of commercial broilers fed with or without probiotic, prebiotic or synbiotic supplemented feed.

Carcass trait	Treatments					CD value
	T ₁	T ₂	T ₃	T ₄	T ₅	
Pre-slaughter weight (g)	1965.00±46.94	1950.00±20.41	1932.50±34.97	1945.00±35.70	1925.00±14.43	NS
Dressed weight (g)	1247.75±31.65	1312.75±15.19	1309.50±30.80	1305.00±42.28	1326.50±10.74	NS
Dressing %	63.51 ^a ±0.99	67.35 ^b ±1.06	67.76 ^b ±1.01	67.06 ^b ±1.24	68.92 ^b ±0.64	3.04
Liver weight (g)	40.75±2.21	42.13±1.08	40.50±1.32	45.25±1.70	46.00±2.16	NS
Heart weight (g)	8.75±0.52	9.50±0.64	7.75±0.85	8.75±0.47	7.88±1.08	NS
Gizzard weight (g)	34.25±1.65	36.38±3.41	37.25±0.62	32.75±3.68	34.38±2.62	NS
Giblet weight (g)	83.75±2.77	88.00±4.14	85.50±0.95	86.75±4.49	88.25±3.78	NS
Giblet %	6.72±0.25	6.70±0.30	6.54±0.11	6.64±0.21	6.65±0.25	NS
Abdominal fat weight (g)	38.63 ^b ±1.65	32.50 ^a ±1.50	36.00 ^{ab} ±1.22	31.75 ^a ±1.18	33.75 ^a ±1.75	4.45
Abdominal fat %	3.10 ^b ±0.15	2.47 ^a ±0.09	2.76 ^{ab} ±0.13	2.45 ^a ±0.16	2.54 ^a ±0.11	0.41
Intestinal length (inches)	79.50±2.59	82.25±1.93	78.00±3.34	80.25±0.85	81.00±2.04	NS
Kidney weight (g)	15.75±1.31	14.00±2.04	15.00±1.87	14.25±1.31	16.50±1.89	NS

*Means within row with different superscript differ significantly (p<0.05), NS=Non-significant

Table-4: Average body weight gain (g) of commercial broilers under different treatments for weekly interval up to 6 weeks age.

Weeks	Treatments					CD value
	T ₁	T ₂	T ₃	T ₄	T ₅	
0-1	81.53 ^a ±1.96	110.10 ^b ±3.62	107.76 ^b ±1.89	103.74 ^b ±3.06	104.45 ^b ±4.45	9.51
1-2	133.98±4.62	152.73±3.90	155.94±4.86	150.91±5.00	155.88±8.19	NS
2-3	296.89±8.15	296.63±9.44	312.30±9.60	313.40±13.05	306.95±9.02	NS
3-4	438.35±10.40	455.50±6.88	461.03±10.84	446.64±12.24	488.32±39.12	NS
4-5	473.18±21.74	477.45±16.47	481.33±9.90	515.35±7.23	488.89±8.68	NS
5-6	489.23±5.21	517.30±28.45	535.58±4.00	499.40±21.42	499.00±17.37	NS
0-4	950.74±12.33	1014.94±17.39	1037.03±12.28	1014.69±30.19	1055.59±39.98	NS
4-6	962.40±25.23	994.75±23.83	1016.90±7.15	1014.75±24.45	987.89±24.60	NS
0-6	1913.14±31.87	2009.69±36.74	2053.93±14.25	2029.44±28.11	2043.49±58.34	NS

*Means within row with different superscript differ significantly (p<0.05)

control groups. The average dressing percentage for different treatment groups is shown in Figure-1.

Liver weight

The highest liver weight was observed in synbiotic half level group (T₅), which was followed by T₄, T₂, T₁ and T₃ group. There were a non-significant difference between all the treatment groups for liver weight [13,14,17,18] found similar findings.

Heart weight

The heart weight with the highest value observed in the probiotic group (T₂) which was followed by T₁, T₄, T₅ and T₃ group. There was non-significant difference between all the treatment groups [13,14,17] found similar findings to this study.

Gizzard weight

The highest gizzard weight was observed in the prebiotic group (T₃) which was followed by T₂, T₅, T₁ and T₄ group. There was non-significant difference between all the treatment groups [13,17,18] also reported similar findings. The average weight of the organs has been shown in Figure-2.

Giblet weight

The highest giblet weight was observed in synbiotic half level group (T₅) which was followed by T₂, T₄, T₃ and T₁ group. There were non-significant differences between all the treatment groups.

Abdominal fat weight and %

The control group (T₁) birds were having highest abdominal fat % which was significantly (p<0.05) higher than T₅, T₂ and T₄ group though there was a non-significant difference between control and prebiotic supplemented group (T₃). The lowest abdominal fat % was observed in the synbiotic group (T₄). The highest abdominal fat % in the control group was followed by T₃, T₅, T₂ and T₄ groups. Present study was in accordance with results of [11,14,19]. Present study differed from results of [12,15,16,20].

Intestinal length

The highest intestinal length was found in the probiotic supplemented group (T₂) which was followed by T₅, T₄, T₁ and T₃. The intestinal length was

not differing among each other though numerical differences were there. Present study was in accordance with findings of [16] but values for intestinal length were lower than present findings.

Kidney weight

The highest kidney weight was noticed in the synbiotic half level group (T₅) which was followed by T₁, T₃, T₄ and T₂ group and there were non-significant differences amongst each other. Present study was in accordance with the results of [14] who observed lower kidney weight than the present study.

Amongst all carcass traits, dressing percentage, abdominal fat weight and abdominal fat percentage (as a percentage of dressed weight) were recorded significant (p<0.05) difference amongst different treatment groups. Synbiotic half level supplemented (T₅) was dominating for most of the traits. Higher dressed weight, higher dressing percentage, higher liver weight, higher giblet weight and higher kidney weight was observed in synbiotic half level supplemented group (T₅) than all other treatments. Synbiotic group (T₄) was having lower abdominal fat weight and lower abdominal fat percentage (as a percentage of dressed weight) among all treatment. Prebiotic group (T₃) had higher gizzard weight than other groups. Whereas, a probiotic group (T₂) had higher heart weight and longer intestinal length amongst all supplemental groups. Giblet percentage (as a percentage of dressed weight) was higher in the control group.

ROFC

Average ROFC in terms of (Rs./bird), (%/bird), income from selling and feed cost of broilers under different treatment groups has been shown in Table-5 and Figure-3. Cost of feed per kg for different treatment groups is shown in Table-6. ROFC in terms of % per bird was indicating the per cent of profit in treatment as compared to the control group. Average daily feed consumption (g/bird/day) is given in Table-7. The income from selling of the birds was significantly (p<0.05) higher in all treatment groups than the control group but there was a non-significant difference between supplemented groups. Feed cost during whole experimental




Figure-1: The average dressing percentages of birds under different treatments.




Figure-2: The average weight of organs included in giblet under different treatment groups.

Table-5: The ROFC (Rs/bird) and (%/bird) realized under different feed supplement groups.

Particulars	Treatment					CD value
	T ₁	T ₂	T ₃	T ₄	T ₅	
Income from bird sold (Rs./bird)	141.32 ^a ±2.25	148.28 ^b ±2.62	151.44 ^b ±1.01	149.67 ^b ±2.03	150.75 ^b ±1.31	5.85
Feed cost (Rs./bird)	85.50 ^b ±1.34	83.63 ^b ±2.38	84.32 ^b ±0.60	78.85 ^a ±1.14	78.47 ^a ±1.18	4.37
ROFC (Rs./bird)	55.82 ^a ±1.13	64.64 ^b ±0.72	67.12 ^{bc} ±1.57	70.82 ^{cd} ±2.38	72.28 ^d ±1.43	4.66
ROFC (Rs./kg live weight)	28.43	31.38	31.91	34.06	34.52	-
ROFC (%/bird)	00.00	15.80	20.24	26.87	29.48	-

ROFC=Return over feed cost, *Means within row with different superscript differ significantly (p<0.05)

Table-6: Cost of feed of control and supplemented with probiotic, prebiotic and synbiotic.

Type of feed	T ₁	T ₂	T ₃	T ₄	T ₅
Broiler starter (Rs./kg)	22.32	22.49	22.39	22.56	22.44
Broiler finisher (Rs./kg)	22.02	22.19	22.09	22.26	22.14



period was significantly (p<0.05) lower in synbiotic supplemented groups (T₄ and T₅) than other groups. ROFC of the control group found significantly (p<0.05) lower than all treatment groups. Non-significant difference was found between T₂ and T₃, T₃ and T₄ and T₄ and T₅. In terms of percentage, highest ROFC was in T₅ (29.48%) than T₄ (26.87%) followed by T₃ (20.24%) and T₂ (15.80%) as compared to control (T₁) suggesting incorporation of synbiotic yields more return and we can afford its 50% level for better economy. The present study was in accordance with [11,21,22] though they observed lower values of profit as compared to present findings, but the trend was same.

EPEI

The EPEI for performance of the broilers is given in Table-8 and Figure-4. The EPEI value was higher in T₄ (285.76) which was followed by T₃ (271.34), T₂ (263.24), T₅ (261.20) and T₁ (231.78). All supplemented groups were having significantly (p<0.05) higher EPEI value than the control. There was a non-significant difference between T₂, T₃, T₅ and T₂, T₃, T₄. The higher EPEI value means higher average body weight; good livability and higher feed efficiency in stipulated number of days thus give overall economics of the birds considering various important traits. Hence, T₄ group (synbiotic) found more economical than other group when EPEI considered.

Conclusion

Dressing percentage was highest in synbiotic (50% level) group (T₅) over other supplemented groups and differed significantly with control in spite of having lowest pre-slaughter weight. The ROFC (Rs.) was significantly (p<0.05) higher in the synbiotic group (T₄ and T₅) than other groups. Also, ROFC in terms of per cent per bird was highest in synbiotic (50% level) group (T₅) than all other groups. EPEI was highest in symbiotic (100%) group (T₄) over other group which means higher average body weight; good livability and higher feed efficiency in stipulated number of days by the group.

**Figure-3:** The Return Over Feed Cost (ROFC) for various treatment groups in commercial broilers.**Figure-4:** European Performance Efficiency Index (EPEI) for various treatment groups in commercial broilers.

Overall it can be concluded that the diet supplemented with synbiotic (100% level) was most efficient in terms of EPEI and synbiotic (50% level) in terms of ROFC. Hence, As feed supplement, synbiotic has a beneficial effect over probiotic and prebiotic when used alone.

Authors' Contributions

RSJ designed the experiment. MAS, FPS and RSJ carried out the research. MAS, ABP and RKM drafted and revised the manuscript. ABP, NJB, RSJ and FPS recorded and analysed the data All authors read and approved the final manuscript.

Acknowledgments

The authors are thankful to the Dean, College of Veterinary Science and Animal Husbandry, Anand,

Table-7: Average daily feed consumption (g/bird/day) of commercial broilers for different treatments during different time intervals.

Weeks	Treatments					CD value
	T ₁	T ₂	T ₃	T ₄	T ₅	
0-1	20.13±0.14	19.31±0.18	19.38±0.20	19.50±0.35	19.16±0.56	NS
1-2	41.58 ^b ±0.50	36.83 ^a ±0.12	36.64 ^a ±0.17	37.08 ^a ±0.30	36.86 ^a ±0.15	0.86
2-3	67.84±0.84	69.92±1.03	69.60±0.43	69.37±0.59	68.59±0.68	NS
3-4	133.26 ^c ±1.26	126.70 ^c ±4.78	129.34 ^c ±2.47	115.48 ^b ±1.57	107.13 ^a ±2.04	8.23
4-5	152.77±4.34	137.70±6.25	136.33±2.00	137.79±4.10	136.75±2.31	NS
5-6	135.54 ^{ab} ±3.95	144.56 ^{bc} ±6.50	150.55 ^c ±0.98	123.58 ^a ±5.34	134.73 ^{ab} ±3.66	13.54
0-4	65.70 ^d ±0.51	63.19 ^c ±1.09	63.74 ^{cd} ±0.67	60.36 ^b ±0.52	57.93 ^a ±0.58	2.14
4-6	144.16±3.95	141.13±5.58	143.44±1.03	130.68±2.97	135.74±2.83	NS
0-6	91.85 ^b ±1.44	89.17 ^b ±2.54	90.31 ^b ±0.64	83.80 ^a ±1.21	83.87 ^a ±1.26	4.67

*Means within row with different superscript differ significantly (p<0.05)

Table-8: The EPEI of broilers fed with Probiotic, Prebiotic and their combination under different treatment.

Particulars	Treatment					CD value
	T ₁	T ₂	T ₃	T ₄	T ₅	
EPEI	231.78 ^a ±4.39	263.24 ^{bc} ±2.99	271.34 ^{bc} ±5.39	285.76 ^c ±9.07	261.20 ^b ±12.17	22.87

EPEI=European performance efficiency index, *Means within row with different superscript differ significantly (p<0.05)

Gujarat for providing necessary funds and facilities for this research.

Competing Interests

The authors declare that they have no competing interests.

References

- Prabhakaran, R. (2012) Overview of poultry production in India vis-a-vis global scenario. In the Proceeding of XXIX Annual Conference and National Symposium of Indian Poultry Science Association (IPSACON-2012) on Commercial and Rural Poultry Production: Novel Concepts and Strategies to Meet Growing Demand and Changing Consumers Needs. 5th-7th December at Hyderabad. p3-11.
- Kokje, R.P. (1999) Effect of feeding probiotics on growth performance in commercial broilers. M. V. Sc. Thesis Submitted to Gujarat Agricultural University, Anand.
- Hinton, M., Kaukas, A. and Linton, A.H. (1986) The ecology of drug resistance in enteric lactobacilli on the performance of broilers. *J. Appl. Bacteriol.*, 15: 77-92. (c. f. Sen, S. Ingale, S.L., Kim, Y.W., Kim, J.S., Kim, K.H., Lohakare, J.D., Kim, E.K., Kim, H.S., Ryu, M.H., Kwon, I.K. and Chae, B.J. (2012). Effect of dietary supplementation of probiotics on performance, caecal morphology and small intestinal morphology of broiler chickens. *Anim. Nutr. Feed Technol.*, 12(1): 1-12).
- Barton, M.D. (2000) Antibiotic use in animal feed and its impact on human health. *Nutr. Res. Rev.*, 13: 279-299.
- FAO. (2001) Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Food and Agriculture Organization of the United Nations and World Health Organization Expert Consultation Report Cordoba/Argentina. p1-34.
- Sekhon, B.S. and Jairath, S. (2010) Prebiotics, probiotics and synbiotics: An overview. *J. Pharm. Educ. Res.*, 1(2): 13-36.
- Ivanovic, S., Baltic, M., Popov-Raljić, J., Pisinov, B., Maslic-Strizak, D., Stojanovic, Z., and Pavlovic, I. (2012) The effect of different probiotics on broiler meat quality. *Afr. J. Microbiol. Res.*, 6(5): 937-943.
- Singh, R.P. (2012) Present status and prospects of egg and poultry processing in India. XXIX annual conference and national symposium of Indian poultry science association (IPSACON-2012) on commercial and rural poultry production: Novel concepts and strategies to meet growing demand and changing consumers needs. 5th-7th December at Hyderabad. p111-115.
- Sharma, N. (2007) Effect of supplementation of enzymes on growth performance in commercial broilers. A Thesis Submitted to Anand Agricultural University. p49.
- Snedecor, G.W. and Cochran, W.G. (1995) Statistical Methods. 8th ed. The Iowa State University Press, Ames, Iowa, USA.
- Anjum, M.I. Khan, A.G. Azim, A. and Afzal, M. (2005) Effect of dietary supplementation of multi-strain probiotic on broiler growth performance. *Pak. Vet. J.*, 25(1): 25-29.
- Bozkurt, M., Kucukayilmaz, K., Catli, A.U. and Cinar, M. (2009) The effect of single or combined dietary supplementation of prebiotics, organic acid and probiotics on performance and slaughter characteristics of broilers. *S. Afr. J. Anim. Sci.*, 39(3): 197-205.
- Abdel-Raheem, M. and Abd-Allah, M.S. (2011) The effect of single or combined dietary supplementation of mannan oligosaccharide and probiotics on performance and slaughter characteristics of broilers. *Int. J. Poult. Sci.*, 10(11): 854-862.
- Sojoudi, M.R., Dadashbeiki, M. and Bouyeh, M. (2012) Effect of different levels of prebiotics TechnoMos on carcass characteristics of broiler chickens. *J. Basic Appl. Sci. Res.*, 2(7): 6778-6794.
- Mohamed, M.A., Hassan, H.M.A. and El-Barkouky, E.M.A. (2008) Effect of mannan oligosaccharide on performance and carcass characteristics of broiler chicks. *J. Agric. Soc. Sci.*, 4(1): 13-17.
- Toghyani, M., Toghyani, M. and Tabeidian, S.A. (2011) Effect of probiotic and prebiotic as antibiotic growth promoter substitutions on productive and carcass traits of broiler chicks. *Int. Conf. Food Eng. Biotechnol.*, 9: 82-86.
- Awad, W.A., Ghareeb, K., Abdel-Raheem, S. and Bohm, J. (2009) Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights and intestinal histomorphology of broiler chickens. *Poult. Sci.*, 88: 49-55.
- Dizaji, B.R. Hejazi, S. and Zakeri, A. (2012) Effects of dietary supplementations of prebiotics, probiotics, synbiotics and acidifiers on growth performance and organs weights of broiler chicken. *Eur. J. Exp. Biol.*, 2(6): 2125-2129.
- Swain, B.K., Naik, P.K., Chakurkar, E.B. and Singh, N.P. (2012) Effect of combined supplementation of probiotic

- and yeast on growth, carcass characteristics and economics of production in broiler chickens. *Anim. Nutr. Feed Technol.*, 12: 103-110.
20. Ashayerizadeh, A., Dabiri, N., Mirzadeh, K. and Ghorbani, M.R. (2011) Effects of dietary inclusion of several biological feed additives on growth response of broiler chickens. *J. Cell Anim. Biol.*, 5(4): 61-65.
21. Ramlah, A.H. and Tan, C.K. (1995) Effects of probiotic supplementation on broiler performance. *Perlanika J. Trop. Agric. Sci.*, 18(2): 109-112.
22. Munj, C.P., Ranade, A.S., Desai, D.N., Patil, M.B., Avari, P.E., Patil, U.M. and Metakari, D.V. (2010) Synergistic effects of feed additives on performance of broilers. *Indian J. Poult. Sci.*, 45(3): 292-296.
